

# (1*H*-Imidazol-4-yl)methanol

Marisa B. Sanders, John C. Farrok, Joseph Hardie and Benny C. Chan\*

Department of Chemistry, The College of New Jersey, 2000 Pennington Rd, Ewing, NJ 08628, USA

Correspondence e-mail: chan@tcnj.edu


Received 5 March 2013; accepted 12 June 2013

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.035;  $wR$  factor = 0.095; data-to-parameter ratio = 17.8.

The title compound,  $\text{C}_4\text{H}_6\text{N}_2\text{O}$ , displays two predominant hydrogen-bonding interactions in the crystal structure. The first is between the unprotonated imidazole N atom of one molecule and the hydroxy H atom of an adjacent molecule. The second is between the hydroxy O atom of one molecule and the imidazole N—H group of a corresponding molecule. These interactions lead to the formation of a two-dimensional network parallel to  $(10\bar{1})$ . C—H $\cdots$ O interactions also occur.

## Related literature

For background information on imidazole complex formation, see: Bauman & Wang (1964); Fan *et al.* (2000). For related structures, see: Nyamori *et al.* (2010); Albov *et al.* (2006). For the use of imidazole-containing compounds in coordination chemistry, see: Huff *et al.* (1993); Fujita *et al.* (1994). For the use of the title compound in the synthesis of biological compounds, see: Darby *et al.* (1942).



## Experimental

### Crystal data

$\text{C}_4\text{H}_6\text{N}_2\text{O}$   
 $M_r = 98.11$   
 Monoclinic,  $C2/c$   
 $a = 13.9180$  (9) Å  
 $b = 7.1980$  (5) Å  
 $c = 11.6509$  (12) Å  
 $\beta = 125.249$  (1)°  
 $V = 953.20$  (13) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation

$\mu = 0.10$  mm<sup>-1</sup>  
 $T = 100$  K

$0.52 \times 0.37 \times 0.29$  mm

### Data collection

Bruker APEXII CCD  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2011)  
 $T_{\min} = 0.688$ ,  $T_{\max} = 0.746$

5389 measured reflections  
 1158 independent reflections  
 1086 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.015$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$   
 $wR(F^2) = 0.095$   
 $S = 1.07$   
 1158 reflections

65 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.35$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.27$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}^{\text{i}}$	0.84	1.92	2.7563 (13)	172
$\text{N2}-\text{H2}\cdots\text{O1}^{\text{ii}}$	0.88	1.99	2.8315 (11)	161
$\text{C4}-\text{H4}\cdots\text{O1}^{\text{iii}}$	0.95	2.57	3.4574 (17)	155

Symmetry codes: (i)  $-x+1, -y, -z+2$ ; (ii)  $-x+\frac{1}{2}, -y+\frac{1}{2}, -z+2$ ; (iii)  $-x+\frac{1}{2}, y+\frac{1}{2}, -z+\frac{3}{2}$ .

Data collection: *APEX2* (Bruker, 2011); cell refinement: *SAINT* (Bruker, 2011); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *CrystalMaker* (CrystalMaker Software, 2009); software used to prepare material for publication: *enCIFer* (Allen *et al.* 2004).

The authors gratefully acknowledge The College of New Jersey's School of Science for research funding and the National Science Foundation for major research instrumentation grant (NSF-0922931) for diffractometer acquisition.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2622).

## References

- Albov, D. V., Rybakov, V. B., Babaev, E. V. & Tsisevich, A. A. (2006). *Acta Cryst.* **E62**, o963–o965.  
 Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.  
 Bauman, J. & Wang, J. (1964). *Inorg. Chem.* **3**, 368–373.  
 Bruker (2011). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 CrystalMaker Software (2009). *CrystalMaker for Windows*. CrystalMaker Software Ltd, Oxford, England.  
 Darby, W., Lewis, H. & Totter, J. (1942). *J. Am. Chem. Soc.* **2**, 463–464.  
 Fan, C., Li, G., Zhu, D. & Xhu, J. (2000). *Chin. J. Chem.* **18**, 115–117.  
 Fujita, M., Kwon, Y. J., Washizu, S. & Ogura, K. (1994). *J. Am. Chem. Soc.* **116**, 1151–1152.  
 Huff, A., Chang, C., Cooper, D., Smith, K. & Dawson, J. (1993). *Inorg. Chem.* **32**, 1460–1466.  
 Nyamori, V. O., Bala, M. D. & Levendis, D. C. (2010). *Acta Cryst.* **E66**, m412.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supplementary materials

*Acta Cryst.* (2013). E69, o1151 [doi:10.1107/S160053681301636X]

**(1*H*-Imidazol-4-yl)methanol**

**Marisa B. Sanders, John C. Farrokh, Joseph Hardie and Benny C. Chan**

**Comment**

Imidazole ligands have been used in coordination chemistry with great success over the last twenty years (Huff *et al.*, 1993). These successes can be attributed to how the nitrogen in imidazole assists in the formation of metal complexes (Fujita *et al.*, 1994). Imidazole-containing metal complexes have a variety of applications, such as redox mediators in enzyme-based electrochemical sensors (Fan *et al.*, 2000). A few examples of imidazole complex compounds with biological applications have been reported (Bauman and Wang, 1964). Histidine, an essential amino acid, and histamine, a bioorganic compound that acts as neurotransmitter, both involve (1*H*-imidazol-5-yl)methanol in their respective syntheses (Darby *et al.*, 1942). Here we report on a new imidazole compound, the hydroxymethyl-substituted imidazole, the title compound, C<sub>4</sub>H<sub>6</sub>N<sub>2</sub>O. The bond lengths and bond angles are within normal ranges in the molecular structure of this compound (Fig. 1). The compound forms hydrogen bonds of 1.985 (8) Å between the nitrogen (N1) on the imidazole ring of one molecule and the hydrogen (H1') of the hydroxyl on an adjacent molecule (Fig. 2). Hydrogen bonding also takes place between the oxygen (O1") on the hydroxyl group of one molecule and the hydrogen (H2) bonded to a nitrogen (N2) on the imidazole ring of a corresponding molecule. This bond measures 1.921 (1) Å (Fig. 3).

**Experimental**


Approximately 100 mg of the target compound was dissolved in 2 ml of a 50% methanol: 50% toluene solution. The solution was allowed to evaporate slowly for two weeks until clear, colorless crystals formed. A crystal was isolated and analyzed on a Bruker *APEX* II CCD single-crystal X-ray diffractometer.

**Refinement**

The structure was solved using direct methods (Bruker, 2011).


**Computing details**

Data collection: *APEX2* (Bruker, 2011); cell refinement: *SAINT* (Bruker, 2011); data reduction: *SAINT* (Bruker, 2011); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *CrystalMaker* (CrystalMaker Software, 2009); software used to prepare material for publication: *enCIFer* (Allen *et al.* 2004).




**Figure 1**

Thermal ellipsoid plot at 50% probability.



**Figure 2**

The title structure is stabilized by hydrogen bonds between N1' and H1, which each measure 1.985 (8) Å. Oxygen atoms are shown in red, carbon atoms in black, hydrogen atoms in pink, and nitrogen atoms in blue.



**Figure 3**

Hydrogen bonding between the oxygen (O1'') on the hydroxyl group of one molecule and the hydrogen (H2) bonded to a nitrogen (N2) on the imidazole ring of a corresponding molecule measures 1.921 (1) Å.

**(1*H*-Imidazol-4-yl)methanol**

*Crystal data*

C<sub>4</sub>H<sub>6</sub>N<sub>2</sub>O

*M<sub>r</sub>* = 98.11

Monoclinic, *C*2/*c*

*a* = 13.9180 (9) Å

*b* = 7.1980 (5) Å

*c* = 11.6509 (12) Å

β = 125.249 (1)°

*V* = 953.20 (13) Å<sup>3</sup>

*Z* = 8

*F*(000) = 416

*D<sub>x</sub>* = 1.367 Mg m<sup>-3</sup>

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 189 reflections

θ = 3.6–28.2°

μ = 0.10 mm<sup>-1</sup>

*T* = 100 K

Blocks, colourless

0.52 × 0.37 × 0.29 mm

*Data collection*

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.3333 pixels mm<sup>-1</sup>

ω and φ scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2011)

*T<sub>min</sub>* = 0.688, *T<sub>max</sub>* = 0.746

5389 measured reflections

1158 independent reflections

1086 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.015

θ<sub>max</sub> = 28.6°, θ<sub>min</sub> = 3.4°

*h* = −18→18

*k* = −9→9

*l* = −15→15

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.035$  $wR(F^2) = 0.095$  $S = 1.07$ 

1158 reflections

65 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0511P)^2 + 0.6749P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.002$  $\Delta\rho_{\max} = 0.35 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of ( $F^2$ ) against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.39678 (6)	0.03753 (10)	1.06291 (7)	0.0178 (2)
H1	0.4528	−0.0374	1.093	0.027*
N1	0.43471 (7)	0.23365 (12)	0.85046 (9)	0.0176 (2)
N2	0.27914 (7)	0.41114 (12)	0.71171 (9)	0.0178 (2)
H2	0.2243	0.4731	0.6368	0.021*
C1	0.43706 (9)	0.22305 (14)	1.06767 (10)	0.0189 (2)
H1A	0.5241	0.2245	1.1246	0.023*
H1B	0.4118	0.3067	1.113	0.023*
C2	0.38850 (8)	0.29121 (13)	0.92295 (10)	0.0160 (2)
C3	0.36594 (9)	0.30971 (14)	0.72413 (11)	0.0178 (2)
H3	0.3766	0.2945	0.6512	0.021*
C4	0.29183 (9)	0.40012 (14)	0.83752 (11)	0.0179 (2)
H4	0.2433	0.4567	0.8605	0.021*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0161 (4)	0.0178 (4)	0.0184 (4)	0.0035 (3)	0.0093 (3)	0.0032 (3)
N1	0.0164 (4)	0.0173 (4)	0.0191 (4)	0.0018 (3)	0.0103 (4)	0.0007 (3)
N2	0.0165 (4)	0.0164 (4)	0.0178 (4)	0.0027 (3)	0.0083 (3)	0.0024 (3)
C1	0.0197 (5)	0.0181 (5)	0.0154 (5)	0.0009 (4)	0.0081 (4)	−0.0012 (4)
C2	0.0159 (5)	0.0140 (4)	0.0169 (5)	−0.0009 (3)	0.0088 (4)	−0.0016 (3)
C3	0.0182 (5)	0.0168 (5)	0.0193 (5)	0.0005 (4)	0.0113 (4)	0.0005 (4)
C4	0.0184 (5)	0.0165 (5)	0.0196 (5)	0.0019 (4)	0.0114 (4)	−0.0001 (4)

*Geometric parameters (Å, °)*

O1—C1	1.4369 (12)	C1—C2	1.4917 (13)
O1—H1	0.84	C1—H1A	0.99
N1—C3	1.3251 (13)	C1—H1B	0.99
N1—C2	1.3877 (12)	C2—C4	1.3674 (14)
N2—C3	1.3459 (13)	C3—H3	0.95
N2—C4	1.3738 (13)	C4—H4	0.95
N2—H2	0.88		
C1—O1—H1		H1A—C1—H1B	
C3—N1—C2		C4—C2—N1	
C3—N2—C4		C4—C2—C1	
C3—N2—H2		N1—C2—C1	
C4—N2—H2		N1—C3—N2	
O1—C1—C2		N1—C3—H3	
O1—C1—H1A		N2—C3—H3	
C2—C1—H1A		C2—C4—N2	
O1—C1—H1B		C2—C4—H4	
C2—C1—H1B		N2—C4—H4	

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O1—H1 $\cdots$ N1 <sup>i</sup>	0.84	1.92	2.7563 (13)	172
N2—H2 $\cdots$ O1 <sup>ii</sup>	0.88	1.99	2.8315 (11)	161
C4—H4 $\cdots$ O1 <sup>iii</sup>	0.95	2.57	3.4574 (17)	155

Symmetry codes: (i)  $-x+1, -y, -z+2$ ; (ii)  $-x+1/2, -y+1/2, -z+2$ ; (iii)  $-x+1/2, y+1/2, -z+3/2$ .